

# NORMS ON COMPLEX MATRICES INDUCED BY COMPLETE HOMOGENEOUS SYMMETRIC POLYNOMIALS

KONRAD AGUILAR, ÁNGEL CHÁVEZ, STEPHAN RAMON GARCIA, AND JURIJ VOLČIČ

**ABSTRACT.** We introduce a remarkable new family of norms on the space of  $n \times n$  complex matrices. These norms arise from the combinatorial properties of symmetric functions, and their construction and validation involve probability theory, partition combinatorics, and trace polynomials in noncommuting variables. Our norms enjoy many desirable analytic and algebraic properties, such as an elegant determinantal interpretation and the ability to distinguish certain graphs that other matrix norms cannot. Furthermore, they give rise to new dimension-independent tracial inequalities. Their potential merits further investigation.

## 1. INTRODUCTION

In this note we introduce a family of norms on complex matrices. These are initially defined in terms of certain symmetric functions of eigenvalues of complex Hermitian matrices. The fact that we deal with eigenvalues, as opposed to their absolute values, is notable. First, it prevents standard machinery, such as the theory of symmetric gauge functions, from applying. Second, the techniques used to establish that we indeed have norms are more complicated than one might expect. For example, combinatorics, probability theory, and Lewis' framework for group invariance in convex matrix analysis each play key roles.

These norms on the Hermitian matrices are of independent interest. They can be computed recursively or directly read from the characteristic polynomial. Moreover, our norms distinguish certain pairs of graphs which the standard norms (operator, Frobenius, Schatten-von Neumann, Ky Fan) cannot distinguish.

Our norms extend in a natural and nontrivial manner to all complex matrices. These extensions of our original norms involve partition combinatorics and trace polynomials in noncommuting variables. A Schur convexity argument permits our norms to be bounded below in terms of the mean eigenvalue of a matrix.

These norms, their unusual construction, and their potential applications suggest a host of open problems. We pose several at the end of the paper.

**1.1. Notation.** Denote by  $\mathbb{N}$ ,  $\mathbb{R}$ , and  $\mathbb{C}$ , respectively, the set of natural numbers, real numbers, and complex numbers. Let  $H_n(\mathbb{C})$  denote the set of  $n \times n$  complex Hermitian matrices and  $M_n(\mathbb{C})$  the set of  $n \times n$  complex matrices. Denote the eigenvalues of  $A \in H_n(\mathbb{C})$  by  $\lambda_1(A) \geq \lambda_2(A) \geq \dots \geq \lambda_n(A)$  and define

$$\lambda(A) = (\lambda_1(A), \lambda_2(A), \dots, \lambda_n(A)) \in \mathbb{R}^n.$$

---

*Key words and phrases.* norm, complete homogeneous symmetric polynomial, partition, trace, positivity, convexity, expectation, complexification, trace polynomial, symmetric tensor power.

Third named author supported by NSF grants DMS-1800123 and DMS-2054002. Fourth named author supported by NSF grant DMS-1954709.

We may use  $\lambda$  and  $\lambda_1, \lambda_2, \dots, \lambda_n$  if the matrix  $A$  is clear from context. Let  $\text{diag}(x_1, x_2, \dots, x_n) \in M_n(\mathbb{C})$  denote the  $n \times n$  diagonal matrix with diagonal entries  $x_1, x_2, \dots, x_n$ , in that order. If  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  is understood from context, we may write  $\text{diag}(\mathbf{x})$  for brevity.

**1.2. Complete homogeneous symmetric polynomials.** The *complete homogeneous symmetric (CHS) polynomial* of degree  $d$  in the  $n$  variables  $x_1, x_2, \dots, x_n$  is

$$h_d(x_1, x_2, \dots, x_n) = \sum_{1 \leq i_1 \leq \dots \leq i_d \leq n} x_{i_1} x_{i_2} \cdots x_{i_d}, \quad (1)$$

the sum of all degree  $d$  monomials in  $x_1, x_2, \dots, x_n$  [28, Sec. 7.5]. For example,

$$\begin{aligned} h_0(x_1, x_2) &= 1, \\ h_1(x_1, x_2) &= x_1 + x_2, \\ h_2(x_1, x_2) &= x_1^2 + x_1 x_2 + x_2^2, \quad \text{and} \\ h_3(x_1, x_2) &= x_1^3 + x_1^2 x_2 + x_1 x_2^2 + x_2^3. \end{aligned}$$

Elementary combinatorics confirms that there are precisely  $\binom{n+d-1}{d}$  summands in the definition (1). We often write  $h_d(\mathbf{x})$ , in which  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ , when the number of variables is clear from context.

For  $d$  even and  $\mathbf{x} \in \mathbb{R}^n$ , Hunter proved that  $h_d(\mathbf{x}) \geq 0$ , with equality if and only if  $\mathbf{x} = \mathbf{0}$  [10]. This is not obvious because some of the summands that comprise  $h_d(\mathbf{x})$  (for  $d$  even) may be negative. Hunter's theorem has been reproved many times; see [2], [4, p. 69 & Thm. 3], [6, Cor. 17], [25, Thm. 2.3], and [29, Thm. 1].

**1.3. Partitions and traces.** A *partition* of  $d \in \mathbb{N}$  is an  $r$ -tuple  $\boldsymbol{\pi} = (\pi_1, \pi_2, \dots, \pi_r) \in \mathbb{N}^r$  such that  $\pi_1 \geq \pi_2 \geq \dots \geq \pi_r$  and  $\pi_1 + \pi_2 + \dots + \pi_r = d$ ; the number of terms  $r$  depends on the partition  $\boldsymbol{\pi}$ . We write  $\boldsymbol{\pi} \vdash d$  if  $\boldsymbol{\pi}$  is a partition of  $d$ .

For  $\boldsymbol{\pi} \vdash d$ , define

$$p_{\boldsymbol{\pi}}(x_1, x_2, \dots, x_n) = p_{\pi_1} p_{\pi_2} \cdots p_{\pi_r},$$

in which  $p_k(x_1, x_2, \dots, x_n) = x_1^k + x_2^k + \dots + x_n^k$ . If the length of  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  is clear from context, we often write  $p_{\boldsymbol{\pi}}(\mathbf{x})$  and  $p_k(\mathbf{x})$ , respectively. Another expression for (1) is

$$h_d(x_1, x_2, \dots, x_n) = \sum_{\boldsymbol{\pi} \vdash d} \frac{p_{\boldsymbol{\pi}}(x_1, x_2, \dots, x_n)}{z_{\boldsymbol{\pi}}}, \quad (2)$$

in which the sum runs over all partitions  $\boldsymbol{\pi} = (\pi_1, \pi_2, \dots, \pi_r)$  of  $d$  and

$$z_{\boldsymbol{\pi}} = \prod_{i \geq 1} i^{m_i} m_i!, \quad (3)$$

where  $m_i$  is the multiplicity of  $i$  in  $\boldsymbol{\pi}$  [28, Prop. 7.7.6]. For example, if  $\boldsymbol{\pi} = (4, 4, 2, 1, 1, 1)$ , then  $z_{\boldsymbol{\pi}} = (1^3 3!)(2^1 1!)(4^2 2!) = 384$  [28, (7.17)].

If  $A \in H_n(\mathbb{C})$  has eigenvalues  $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_n)$ , then

$$p_{\boldsymbol{\pi}}(\boldsymbol{\lambda}) = p_{\pi_1}(\boldsymbol{\lambda}) p_{\pi_2}(\boldsymbol{\lambda}) \cdots p_{\pi_r}(\boldsymbol{\lambda}) = (\text{tr } A^{\pi_1})(\text{tr } A^{\pi_2}) \cdots (\text{tr } A^{\pi_r}). \quad (4)$$

This connects eigenvalues, traces, and partitions to symmetric polynomials.

**1.4. Main results.** The following theorem provides a family of novel norms on the space  $H_n(\mathbb{C})$  of  $n \times n$  Hermitian matrices. Some special properties of these norms are discussed in Section 4.

**Theorem 1.** *For even  $d \geq 2$ , the following is a norm on  $H_n(\mathbb{C})$ :*

$$\|A\|_d = (h_d(\lambda_1(A), \lambda_2(A), \dots, \lambda_n(A)))^{1/d}.$$

For example, equations (2) and (4) yield trace-polynomial representations

$$\begin{aligned} \|A\|_2^2 &= \frac{1}{2}(\operatorname{tr}(A^2) + (\operatorname{tr} A)^2), \\ \|A\|_4^4 &= \frac{1}{24}((\operatorname{tr} A)^4 + 6(\operatorname{tr} A)^2 \operatorname{tr}(A^2) + 3(\operatorname{tr}(A^2))^2 + 8(\operatorname{tr} A) \operatorname{tr}(A^3) + 6 \operatorname{tr}(A^4)). \end{aligned} \quad (5)$$

Theorem 1 is nontrivial for several reasons.

- (a) The sums (1) and (2) that characterize  $h_d(\lambda(A))$  may contain negative summands. For example,  $(\operatorname{tr} A) \operatorname{tr}(A^3)$  in (5) can be negative for Hermitian  $A$ .
- (b) The relationship between the spectra of (Hermitian)  $A$ ,  $B$ , and  $A + B$ , conjectured by A. Horn in 1962 [9], was only established in 1998-9 by Klyachko [13] and Knutson–Tao [14]. Therefore, the triangle inequality is difficult to establish. Even if  $A$  and  $B$  are diagonal, the result is not obvious; see (11).
- (c) The sums that define these norms do not involve the absolute values of the eigenvalues of  $A$ . Theorem 1 does not follow from standard considerations, but rather from delicate properties of multivariate symmetric polynomials.

**Example 2.** Because CHS norms do not rely upon the absolute values of the eigenvalues of a Hermitian matrix (that is, its singular values), they can sometimes distinguish singularly cospectral graphs (graphs with the same singular values) that are not cospectral. This feature is not enjoyed by many standard norms (e.g., operator, Frobenius, Schatten–von Neumann, Ky Fan). For example,

$$K = \begin{bmatrix} 0 & 1 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 0 \end{bmatrix},$$

which has eigenvalues  $2, -1, -1$ , is the adjacency matrix for the complete graph on three vertices. The graphs with adjacency matrices

$$A = \begin{bmatrix} K & 0 \\ 0 & K \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & K \\ K & 0 \end{bmatrix}$$

are singularly cospectral but not cospectral: their eigenvalues are  $-1, -1, -1, -1, 2, 2$  and  $-2, -1, -1, 1, 1, 2$ , respectively. However,  $\|A\|_6^6 = 120 \neq 112 = \|B\|_6^6$ .

The norms of Theorem 1 extend in a natural and nontrivial fashion to the set  $M_n(\mathbb{C})$  of all  $n \times n$  complex matrices.

**Theorem 3.** *Let  $d \geq 2$  be even and let  $\pi = (\pi_1, \pi_2, \dots, \pi_r)$  be a partition of  $d$ . Define  $T_\pi : M_n(\mathbb{C}) \rightarrow \mathbb{R}$  by setting  $T_\pi(A)$  to be  $1/\binom{d}{d/2}$  times the sum over the  $\binom{d}{d/2}$  possible locations to place  $d/2$  adjoints\* among the  $d$  copies of  $A$  in*

$$(\operatorname{tr} \underbrace{AA \cdots A}_{\pi_1}) (\operatorname{tr} \underbrace{AA \cdots A}_{\pi_2}) \cdots (\operatorname{tr} \underbrace{AA \cdots A}_{\pi_r}).$$

Then

$$\|A\|_d = \left( \sum_{\pi} \frac{T_{\pi}(A)}{z_{\pi}} \right)^{1/d}, \quad (6)$$

in which the sum runs over all partitions  $\pi$  of  $d$  and  $z_{\pi}$  is defined in (3), is a norm on  $M_n(\mathbb{C})$  that restricts to the norm on  $H_n(\mathbb{C})$  given by Theorem 1.

**Example 4.** The two partitions of  $d = 2$  satisfy  $z_{(2)} = 2$  and  $z_{(1,1)} = 2$ . There are  $\binom{2}{1} = 2$  ways to place two adjoints  $*$  in a string of two  $A$ s. Therefore,

$$T_{(2)}(A) = \frac{1}{2}(\operatorname{tr}(A^*A) + \operatorname{tr}(AA^*)) = \operatorname{tr}(A^*A), \quad \text{and}$$

$$T_{(1,1)}(A) = \frac{1}{2}((\operatorname{tr} A^*)(\operatorname{tr} A) + (\operatorname{tr} A)(\operatorname{tr} A^*)) = (\operatorname{tr} A^*)(\operatorname{tr} A),$$

so

$$\|A\|_2^2 = \frac{1}{2} \operatorname{tr}(A^*A) + \frac{1}{2} (\operatorname{tr} A^*)(\operatorname{tr} A). \quad (7)$$

**Example 5.** The five partitions of  $d = 4$  satisfy  $z_{(4)} = 4$ ,  $z_{(3,1)} = 3$ ,  $z_{(2,2)} = 8$ ,  $z_{(2,1,1)} = 4$ , and  $z_{(1,1,1,1)} = 24$ . There are  $\binom{4}{2} = 6$  ways to place two adjoints  $*$  in a string of four  $A$ s. For example,

$$\begin{aligned} 6T_{(3,1)}(A) &= (\operatorname{tr} A^*A^*A)(\operatorname{tr} A) + (\operatorname{tr} A^*AA^*)(\operatorname{tr} A) + (\operatorname{tr} A^*AA)(\operatorname{tr} A^*) \\ &\quad + (\operatorname{tr} AA^*A^*)(\operatorname{tr} A) + (\operatorname{tr} AA^*A)(\operatorname{tr} A^*) + (\operatorname{tr} AAA^*)(\operatorname{tr} A^*) \\ &= 3 \operatorname{tr}(A^{*2}A)(\operatorname{tr} A) + 3(\operatorname{tr} A^2A^*)(\operatorname{tr} A^*). \end{aligned}$$

Some computation and (6) reveal that

$$\begin{aligned} \|A\|_4^4 &= \frac{1}{24} ((\operatorname{tr} A)^2 \operatorname{tr}(A^*)^2 + \operatorname{tr}(A^*)^2 \operatorname{tr}(A^2) + 4 \operatorname{tr}(A) \operatorname{tr}(A^*) \operatorname{tr}(A^*A) \\ &\quad + 2 \operatorname{tr}(A^*A)^2 + (\operatorname{tr} A)^2 \operatorname{tr}(A^{*2}) + \operatorname{tr}(A^2) \operatorname{tr}(A^{*2}) + 4 \operatorname{tr}(A^*) \operatorname{tr}(A^*A^2) \\ &\quad + 4 \operatorname{tr}(A) \operatorname{tr}(A^{*2}A) + 2 \operatorname{tr}(A^*AA^*A) + 4 \operatorname{tr}(A^{*2}A^2)). \end{aligned} \quad (8)$$

If  $A = A^*$ , this simplifies to the norm (5) on  $H_n(\mathbb{C})$ , as expected.

Because of their origins in terms of complete homogeneous symmetric polynomials, we sometimes refer to the norm  $\|\cdot\|_d$  as the *CHS norm* of order  $d$ . The notation  $\|\cdot\|$  is used, occasionally with subscripts, for other norms.

In the Hermitian case, the norm  $\|\cdot\|_d$  can be directly extracted from the Taylor expansion of an explicit rational function (Theorem 19). The general situation is elegantly summarized in a determinantal formula.

**Theorem 6.** Let  $A \in M_n(\mathbb{C})$ . For  $d$  even,  $\binom{d}{d/2} \|A\|_d^d$  is the coefficient of  $z^{d/2} \bar{z}^{d/2}$  in the Taylor expansion of  $\det(I - zA - \bar{z}A^*)^{-1}$  about the origin.

Helton and Vinnikov showed that polynomials of the form  $p = \det(I - zA - \bar{z}A^*) \in \mathbb{C}[z, \bar{z}]$  are precisely the real-zero polynomials in  $\mathbb{C}[z, \bar{z}]$  [8]. That is, they are characterized by the conditions  $p(0) = 1$  and that  $x \mapsto p(\alpha x)$  has only real zeros for every  $\alpha \in \mathbb{C}$ . Properties of such polynomials are studied within the framework of hyperbolic [23] and stable [32] polynomials.

This paper is structured as follows. Sections 2 and 3 contain the proofs of Theorems 1 and 3, respectively. Section 4 surveys the remarkable properties of the CHS norms, such as Theorem 6. We pose several open questions in Section 5.

2. PROOF OF THEOREM 1

Let  $d \geq 2$  be even. We prove that  $\mathfrak{H} : H_n(\mathbb{C}) \rightarrow \mathbb{R}$  defined by

$$\mathfrak{H}(A) = h_d(\lambda_1(A), \lambda_2(A), \dots, \lambda_n(A))^{1/d} \quad (9)$$

is a norm. Hunter's theorem ensures that  $\mathfrak{H}(A) \geq 0$  and, moreover, that  $\mathfrak{H}(A) = 0$  if and only if  $A = 0$  (in fact, the nonnegativity of  $\mathfrak{H}$  follows from (10) below). Since  $\mathfrak{H}(cA) = |c|\mathfrak{H}(A)$  for all  $A \in H_n(\mathbb{C})$  and  $c \in \mathbb{R}$ , it suffices to prove that  $\mathfrak{H}$  satisfies the triangle inequality. This is accomplished by combining Lewis' framework for group invariance in convex matrix analysis [16] with a probabilistic approach to the complete homogeneous symmetric polynomials [1, 25, 29].

**2.1. Group invariance.** Let  $\mathcal{V}$  be a finite-dimensional  $\mathbb{R}$ -inner product space. The adjoint  $\varphi^*$  of a linear map  $\varphi : \mathcal{V} \rightarrow \mathcal{V}$  satisfies  $\langle \varphi^*(X), Y \rangle = \langle X, \varphi(Y) \rangle$  for all  $X, Y \in \mathcal{V}$ . We say that  $\varphi$  is *orthogonal* if  $\varphi^* \circ \varphi$  is the identity map on  $\mathcal{V}$ . Let  $O(\mathcal{V})$  denote the set of all orthogonal linear maps on  $\mathcal{V}$ . For a subgroup  $G \subseteq O(\mathcal{V})$ , we say that  $f : \mathcal{V} \rightarrow \mathbb{R}$  is *G-invariant* if  $f(\varphi(X)) = f(X)$  for all  $\varphi \in G$  and  $X \in \mathcal{V}$ .

**Definition 7** (Def. 2.1 of [16]).  $\delta : \mathcal{V} \rightarrow \mathcal{V}$  is a *G-invariant normal form* if

- (a)  $\delta$  is G-invariant,
- (b) For each  $X \in \mathcal{V}$ , there is an  $\varphi \in O(\mathcal{V})$  such that  $X = \varphi(\delta(X))$ , and
- (c)  $\langle X, Y \rangle \leq \langle \delta(X), \delta(Y) \rangle$  for all  $X, Y \in \mathcal{V}$ .

In this case,  $(\mathcal{V}, G, \delta)$  is a *normal decomposition system*.

Suppose that  $(\mathcal{V}, G, \delta)$  is a normal decomposition system and  $\mathcal{W} \subseteq \mathcal{V}$  is a subspace. The *stabilizer* of  $\mathcal{W}$  in  $G$  is  $G_{\mathcal{W}} = \{\varphi \in G : \varphi(\mathcal{W}) = \mathcal{W}\}$ . For convenience, we restrict the domain of each  $\varphi \in G_{\mathcal{W}}$  and consider  $G_{\mathcal{W}}$  as a subset of  $O(\mathcal{W})$ .

Our interest in this material stems from the next result.

**Lemma 8** (Thm. 4.3 of [16]). *Let  $(\mathcal{V}, G, \delta)$  and  $(\mathcal{W}, G_{\mathcal{W}}, \delta|_{\mathcal{W}})$  be normal decomposition systems with  $\text{ran } \delta \subseteq \mathcal{W}$ . Then a G-invariant function  $f : \mathcal{V} \rightarrow \mathbb{R}$  is convex if and only if its restriction to  $\mathcal{W}$  is convex.*

Let  $\mathcal{V} = H_n(\mathbb{C})$  denote the  $\mathbb{R}$ -inner product space of complex Hermitian matrices, endowed with the inner product  $\langle X, Y \rangle = \text{tr}(XY)$ , and let  $U_n(\mathbb{C})$  denote the group of  $n \times n$  unitary matrices; see Remark 10 for more details about this inner product. For each  $U \in U_n(\mathbb{C})$ , define a linear map  $\varphi_U : \mathcal{V} \rightarrow \mathcal{V}$  by  $\varphi_U(X) = UXU^*$ . Observe that  $\varphi_U \circ \varphi_V = \varphi_{UV}$  and hence

$$G = \{\varphi_U : U \in U_n(\mathbb{C})\}$$

is a group under composition. Since  $\varphi_U^* = \varphi_{U^*}$ , we conclude that  $G$  is a subgroup of  $O(\mathcal{V})$ . Moreover, the function (9) is G-invariant.

Let  $\mathcal{W} = D_n(\mathbb{R})$  denote the set of real diagonal matrices. Then  $\mathcal{W}$  inherits an inner product from  $\mathcal{V}$  and  $G_{\mathcal{W}} = \{\varphi_P : P \in P_n\}$ , in which  $P_n$  denotes the set of  $n \times n$  permutation matrices. Define  $\delta : \mathcal{V} \rightarrow \mathcal{V}$  by

$$\delta(X) = \text{diag}(\lambda_1(X), \lambda_2(X), \dots, \lambda_n(X)),$$

the  $n \times n$  diagonal matrix with  $\lambda_1(X), \lambda_2(X), \dots, \lambda_n(X)$  on its diagonal. Observe that  $\text{ran } \delta \subseteq \mathcal{W}$  since the eigenvalues of a Hermitian matrix are real (in fact,  $\text{ran } \delta = \mathcal{W}$ ). We maintain all of this notation below.

**Lemma 9.**  $(\mathcal{V}, G, \delta)$  and  $(\mathcal{W}, G_{\mathcal{W}}, \delta|_{\mathcal{W}})$  are normal decomposition systems.

*Proof.* We first show that  $(\mathcal{V}, G, \delta)$  is a normal decomposition system. (a) Since eigenvalues are invariant under similarity,  $\delta$  is  $G$ -invariant. (b) For  $X \in \mathcal{V}$ , the spectral theorem provides a  $U \in U_n(\mathbb{C})$  such that  $X = U\delta(X)U^* = \varphi_U(\delta(A))$ . (c) For  $X, Y \in \mathcal{V}$ , note that  $\text{tr } XY \leq \text{tr } \delta(X)\delta(Y)$  [15, Thm. 2.2]; see Remark 10.

We now show that  $(\mathcal{W}, G_{\mathcal{W}}, \delta|_{\mathcal{W}})$  is a normal decomposition system. (a)  $\delta|_{\mathcal{W}}$  is  $G_{\mathcal{W}}$ -invariant since  $\delta(\varphi_P(X)) = \delta(PXP^*) = \delta(X)$  for every  $X \in \mathcal{W}$  and  $P \in P_n$ . (b) Let  $X \in \mathcal{W}$ . Since  $X$  is diagonal there exists a  $P \in P_n$  such that  $X = P\delta(X)P^* = \varphi_P(\delta(X))$ . (c) The diagonal elements of a diagonal matrix are its eigenvalues. Consequently, this property is inherited from  $\mathcal{V}$ ; see Remark 11.  $\square$

**2.2. CHS polynomials as expectations.** Let  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$  be a vector of independent standard exponential random variables [3, (20.10)], and let  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ . Since  $\mathbb{E}[\xi_i^k] = k!$  for  $i = 1, 2, \dots, n$  [3, Ex. 21.3], we deduce that

$$\begin{aligned} \mathbb{E}[|\langle \xi, \mathbf{x} \rangle|^d] &= \mathbb{E}[(\xi_1 x_1 + \xi_2 x_2 \cdots + \xi_n x_n)^d] \\ &= \mathbb{E} \left[ \sum_{k_1+k_2+\dots+k_n=d} \frac{d!}{k_1! k_2! \cdots k_n!} \xi_1^{k_1} \xi_2^{k_2} \cdots \xi_n^{k_n} x_1^{k_1} x_2^{k_2} \cdots x_n^{k_n} \right] \\ &= \sum_{k_1+k_2+\dots+k_n=d} \frac{d!}{k_1! k_2! \cdots k_n!} \mathbb{E} \left[ \xi_1^{k_1} \xi_2^{k_2} \cdots \xi_n^{k_n} x_1^{k_1} x_2^{k_2} \cdots x_n^{k_n} \right] \\ &= d! \sum_{k_1+k_2+\dots+k_n=d} \frac{\mathbb{E}[\xi_1^{k_1}] \mathbb{E}[\xi_2^{k_2}] \cdots \mathbb{E}[\xi_n^{k_n}]}{k_1! k_2! \cdots k_n!} x_1^{k_1} x_2^{k_2} \cdots x_n^{k_n} \\ &= d! \sum_{k_1+k_2+\dots+k_n=d} x_1^{k_1} x_2^{k_2} \cdots x_n^{k_n} \\ &= d! h_d(\mathbf{x}) \end{aligned}$$

for integral  $d \geq 1$  by the linearity of expectation and the independence of the  $\xi_1, \xi_2, \dots, \xi_n$ ; see Remark 12. Now suppose that  $d$  is even. Then

$$h_d(\mathbf{x}) = \mathbb{E}[|\langle \xi, \mathbf{x} \rangle|^d] \geq 0. \quad (10)$$

For  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ , Minkowski's inequality implies that

$$\left( \mathbb{E}[|\langle \xi, \mathbf{x} + \mathbf{y} \rangle|^d] \right)^{1/d} \leq \left( \mathbb{E}[|\langle \xi, \mathbf{x} \rangle|^d] \right)^{1/d} + \left( \mathbb{E}[|\langle \xi, \mathbf{y} \rangle|^d] \right)^{1/d},$$

and hence (for  $d$  even)

$$[h_d(\mathbf{x} + \mathbf{y})]^{1/d} \leq [h_d(\mathbf{x})]^{1/d} + [h_d(\mathbf{y})]^{1/d}. \quad (11)$$

**2.3. Conclusion.** Recall the definition (9) of the function  $\mathfrak{H} : H_n(\mathbb{C}) \rightarrow \mathbb{R}$ . The inequality (11) ensures that the restriction of  $\mathfrak{H}$  to  $D_n(\mathbb{R})$  satisfies the triangle inequality. For  $A, B \in D_n(\mathbb{R})$  and  $t \in [0, 1]$ , note that

$$\mathfrak{H}(tA + (1-t)B) \leq \mathfrak{H}(tA) + \mathfrak{H}((1-t)B) = t\mathfrak{H}(A) + (1-t)\mathfrak{H}(B)$$

by (11) and homogeneity. Thus,  $\mathfrak{H}$  is a convex function on  $D_n(\mathbb{R})$ . Since  $\mathfrak{H}$  is  $G$ -invariant, we conclude from Lemma 8 that  $\mathfrak{H}$  is convex on  $H_n(\mathbb{C})$ . It satisfies the triangle inequality on  $H_n(\mathbb{C})$  since it is convex and homogeneous:

$$\frac{1}{2}\mathfrak{H}(A + B) = \mathfrak{H}\left(\frac{1}{2}A + \frac{1}{2}B\right) \leq \frac{1}{2}\mathfrak{H}(A) + \frac{1}{2}\mathfrak{H}(B).$$

Consequently,  $\mathfrak{H}(A)$  is a norm on  $H_n(\mathbb{C})$ .  $\square$

**2.4. Remarks.** We collect here a few remarks about the proof of Theorem 1.

**Remark 10.** Consider the inner product  $\langle X, Y \rangle = \text{tr}(XY)$  on  $H_n(\mathbb{C})$ ; it is the restriction of the Frobenius inner product to  $H_n(\mathbb{C})$ . The inequality

$$\text{tr}(XY) \leq \text{tr} \delta(X)\delta(Y) \quad \text{for } X, Y \in H_n(\mathbb{C}) \quad (12)$$

is due to von Neumann [31] and has been reproved many times; see de Sá [5], Lewis [15, Thm. 2.2], Marcus [17], Marshall [18], Mirsky [20, Thm. 1], Richter [24, Satz. 1], Rendl and Wolkowicz [22, Cor. 3.1], and Theobald [30].

**Remark 11.** For diagonal matrices, the inequality (12) is equivalent to a classical rearrangement result:  $\langle \mathbf{x}, \mathbf{y} \rangle \leq \langle \tilde{\mathbf{x}}, \tilde{\mathbf{y}} \rangle$ , in which where  $\tilde{\mathbf{x}} \in \mathbb{R}^n$  has the components of  $\mathbf{x} = (x_1, x_2, \dots, x_n)$  in decreasing order [7, Thm. 368].

**Remark 12.** For even  $d$ , (10) implies the nonnegativity of the CHS polynomials. This probabilistic approach appears in the comments on the blog entry [29], and in [27, Lem. 12], which cites [1]. There are many other proofs of the nonnegativity of the even-degree CHS polynomials. Of course, there is Hunter’s inductive proof [10]. Roventa and Temereanca used divided differences [25, Thm. 3.5]. Recently, Bottcher, Garcia, Omar and O’Neill [4] employed a spline-based approach suggested by Olshansky after Garcia, Omar, O’Neill, and Yih obtained it as a byproduct of investigations into numerical semigroups [6, Cor. 17].

**Remark 13.** We stress that our inequality (11) permits  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ ; that is, with no positivity assumptions. For  $p \in \mathbb{N}$ , the similar inequality

$$h_p(\mathbf{x} + \mathbf{y})^{1/p} \leq h_p(\mathbf{x})^{1/p} + h_p(\mathbf{y})^{1/p} \quad \text{for } \mathbf{x}, \mathbf{y} \in \mathbb{R}_{\geq 0}^n \quad (13)$$

has been rediscovered several times. According to McLeod [19, p. 211] and Whiteley [33, p. 49], it was first conjectured by A.C. Aitken. Priority must be given to Whiteley [33, eq. (5)], whose paper appeared in 1958. McLeod’s paper was received on March 16, 1959, although he was unaware of Whiteley’s proof: “To the best of my knowledge, no proof of [(13)] exists so far in the literature.” For more exotic inequalities along the lines of (13), see [27].

### 3. PROOF OF THEOREM 3

The first step in the proof of Theorem 3 is a general complexification result. Let  $\mathcal{V}$  be a complex vector space with a conjugate-linear involution  $v \mapsto v^*$ . Let  $\|\cdot\|$  be a norm on the real subspace  $\mathcal{V}_{\mathbb{R}} = \{v \in \mathcal{V} : v = v^*\}$  of  $*$ -fixed points. For each  $v \in \mathcal{V}$  and  $t \in \mathbb{R}$ , we have  $e^{it}v + e^{-it}v^* \in \mathcal{V}_{\mathbb{R}}$ . Note that the path  $t \mapsto \|e^{it}v + e^{-it}v^*\|$  is continuous for each  $v \in \mathcal{V}$ .

**Proposition 14.** *For even  $d \geq 2$ , the following is a norm on  $\mathcal{V}$  that extends  $\|\cdot\|$ :*

$$\mathfrak{N}_d(v) = \left( \frac{1}{2\pi \binom{d}{d/2}} \int_0^{2\pi} \|e^{it}v + e^{-it}v^*\|^d dt \right)^{1/d}. \quad (14)$$

*Proof.* If  $v \in \mathcal{V}_{\mathbb{R}}$ , then  $\|e^{it}v + e^{-it}v^*\| = |2 \cos t| \|v\|$ . Moreover,  $\mathfrak{N}_d(v) = \|v\|$  since

$$\int_0^{2\pi} |2 \cos t|^d dt = 2\pi \binom{d}{d/2}.$$

Next we verify that  $\mathfrak{N}_d$  is a norm on  $\mathcal{V}$ .

**POSITIVE DEFINITENESS.** The nonnegativity of  $\|\cdot\|$  on  $\mathcal{V}_{\mathbb{R}}$  and (14) ensure that  $\mathfrak{N}_d$  is nonnegative on  $\mathcal{V}$ . If  $v \in \mathcal{V} \setminus \{0\}$ , then  $v = \frac{1}{2}(v + v^*) + i\frac{1}{2}(-iv + iv^*)$ , in which  $v + v^*$  and  $-iv + iv^*$  belong to  $\mathcal{V}_{\mathbb{R}}$ . Thus,  $\|v + v^*\| \neq 0$  or  $\|-iv + iv^*\| \neq 0$ . Since  $t \mapsto \|e^{it}v + e^{-it}v^*\|$  is continuous and nonnegative, it follows that  $\mathfrak{N}_d(v) \neq 0$ .

**ABSOLUTE HOMOGENEITY.** For  $r > 0$  and  $\theta \in \mathbb{R}$ , we have  $\mathfrak{N}_d((re^{i\theta})v) = r\mathfrak{N}_d(e^{i\theta}v) = r\mathfrak{N}_d(v)$  by the  $\mathbb{R}$ -homogeneity of  $\|\cdot\|$  and the periodicity of the integrand in (14).

**TRIANGLE INEQUALITY.** For  $u, v \in \mathcal{V}$ ,

$$\begin{aligned} & \left( \int_0^{2\pi} \|e^{it}(u+v) + e^{-it}(u+v)^*\|^d dt \right)^{1/d} \\ & \leq \left( \int_0^{2\pi} (\|e^{it}u + e^{-it}u^*\| + \|e^{it}v + e^{-it}v^*\|)^d dt \right)^{1/d} \\ & \leq \left( \int_0^{2\pi} \|e^{it}u + e^{-it}u^*\|^d dt \right)^{1/d} + \left( \int_0^{2\pi} \|e^{it}v + e^{-it}v^*\|^d dt \right)^{1/d}, \end{aligned}$$

where the first inequality holds by monotonicity of power functions and the triangle inequality for  $\|\cdot\|$ , and the second inequality holds by the triangle inequality for the  $L^d$  norm on the space  $C[0, 2\pi]$ .  $\square$

There are several natural complexifications of a real Banach space [21]. However, the extensions  $\mathfrak{N}_d$  preserve some of the analytic and algebraic properties of the original norm. For example, if the extension  $\mathfrak{N}_d$  is applied to the norm  $\|\cdot\|_d$  on  $H_n(\mathbb{C})$ , one obtains a norm on  $M_n(\mathbb{C})$  that is given by a trace polynomial.

Let  $\langle x, x^* \rangle$  be the free monoid generated by  $x$  and  $x^*$ . Let  $|w|$  denote the length of a word  $w \in \langle x, x^* \rangle$  and let  $|w|_x$  count the occurrences of  $x$  in  $w$ . For  $A \in M_n(\mathbb{C})$ , let  $w(A) \in M_n(\mathbb{C})$  be the natural evaluation of  $w$  at  $A$ . For example, if  $w = xx^*x^2$ , then  $|w| = 4$ ,  $|w|_x = 3$ , and  $w(A) = AA^*A^2$ .

**Lemma 15.** *Let  $d \geq 2$  be even and let  $\pi = (\pi_1, \dots, \pi_r)$  be a partition of  $d$ . If  $A \in M_n(\mathbb{C})$ , then*

$$\begin{aligned} & \frac{1}{2\pi} \int_0^{2\pi} \text{tr}(e^{it}A + e^{-it}A^*)^{\pi_1} \cdots \text{tr}(e^{it}A + e^{-it}A^*)^{\pi_r} dt \\ & = \sum_{\substack{w_1, \dots, w_r \in \langle x, x^* \rangle: \\ |w_j| = \pi_j \ \forall j \\ |w_1 \cdots w_r|_x = \frac{d}{2}}} \text{tr } w_1(A) \cdots \text{tr } w_r(A). \end{aligned} \quad (15)$$

*Proof.* For every Laurent polynomial  $f \in \mathbb{C}[z, z^{-1}]$  with the constant term  $f_0$  we have  $\int_0^{2\pi} f(e^{it}) dt = 2\pi f_0$ . Let us view

$$f = \text{tr}(zA + z^{-1}A^*)^{\pi_1} \cdots \text{tr}(zA + z^{-1}A^*)^{\pi_r}$$

as a Laurent polynomial in  $z$ . Its constant term is

$$f_0 = \sum_{w_1, \dots, w_r} \text{tr } w_1(A) \cdots \text{tr } w_r(A)$$

where the sum runs over all words  $w_1, w_2, \dots, w_r$  in  $\langle x, x^* \rangle$  with  $|w_j| = \pi_j$  such that the number of occurrences of  $x$  in  $w_1 w_2 \cdots w_r$  equals the number of occurrences of  $x^*$  in  $w_1 w_2 \cdots w_r$ . Thus, (15) follows.  $\square$



Given a partition  $\pi = (\pi_1, \dots, \pi_r)$  of  $d$  and  $A \in M_n(\mathbb{C})$  let

$$T_\pi(A) = \frac{1}{\binom{d}{d/2}} \sum_{\substack{w_1, \dots, w_r \in \langle x, x^* \rangle: \\ |w_j| = \pi_j \ \forall j \\ |w_1 \cdots w_r|_x = \frac{d}{2}}} \text{tr } w_1(A) \cdots \text{tr } w_r(A).$$

We now complete the proof of Theorem 3. The conjugate transpose  $A \mapsto A^*$  is a real structure on  $M_n(\mathbb{C})$ . The corresponding real subspace of  $*$ -fixed points is  $H_n(\mathbb{C})$ . We apply Proposition 14 to the norm  $\|\cdot\|_d$  on  $H_n(\mathbb{C})$  and obtain its extension  $\mathfrak{N}_d(\cdot)$  to  $M_n(\mathbb{C})$  defined by (14). The fact that  $\mathfrak{N}_d(A)$  admits a trace-polynomial expression as in (6) follows from (2) and Lemma 15.  $\square$

**Remark 16.** Proving that (6) is a norm relies crucially on Theorem 1, which states that its restriction to  $H_n(\mathbb{C})$  is a norm. On the other hand, demonstrating that (6) is a norm in a direct manner seems arduous. To a certain degree, this mirrors the current absence of general certificates for dimension-independent positivity of trace polynomials in  $x, x^*$  (see [12] for the analysis in a dimension-fixed setting).

**Remark 17.** For any  $A \in M_n(\mathbb{C})$  and  $t \in [0, 2\pi]$ , the matrices  $e^{it}A + e^{-it}A^*$  are Hermitian. Thus,  $\|e^{it}A + e^{-it}A^*\|_d$  can be defined as in Theorem 1 and hence

$$\|A\|_d = \left( \frac{1}{2\pi \binom{d}{d/2}} \int_0^{2\pi} \|e^{it}A + e^{-it}A^*\|_d^d dt \right)^{1/d}. \quad (16)$$

**Remark 18.** Here is another way to restrict  $\|\cdot\|_d$  to the Hermitian matrices. The proof of Lemma 15 shows that  $\binom{d}{d/2} \|A\|_d^d$  is the coefficient of  $z^{d/2} \bar{z}^{d/2}$  in

$$\|zA + \bar{z}A^*\|_d^d \in \mathbb{C}[z, \bar{z}].$$

#### 4. PROPERTIES OF CHS NORMS

We now establish several properties of the CHS norms. First, we show how the CHS norm of a Hermitian matrix can be computed rapidly and exactly from its characteristic polynomial and recursion (Subsection 4.1). This leads quickly to the determinantal interpretation presented in the introduction (Subsection 4.2). Next, we identify those CHS norms induced by inner products (Subsection 4.3). In Subsection 4.4, we use Schur convexity to provide a lower bound on the CHS norms in terms of the trace seminorm on  $M_n(\mathbb{C})$ . We discuss monotonicity properties in Subsection 4.5 and symmetric tensor powers in Subsection 4.6.

**4.1. Exact computation via characteristic polynomial.** The CHS norm of a Hermitian matrix can be exactly computed from its characteristic polynomial. The following theorem involves only formal series manipulations.

**Theorem 19.** *Let  $p_A(x)$  denote the characteristic polynomial of  $A \in H_n(\mathbb{C})$ . For  $d \geq 2$  even,  $\|A\|_d^d$  is the  $d$ th coefficient in the Taylor expansion of*

$$\frac{1}{\det(I - xA)} = \frac{1}{x^n p_A(1/x)}$$

*about the origin.*

*Proof.* Let  $p_A(x) = (x - \lambda_1)(x - \lambda_2) \cdots (x - \lambda_n)$ . For  $|x|$  small, [29, (1)] provides

$$\sum_{d=0}^{\infty} h_d(\lambda_1, \lambda_2, \dots, \lambda_n) x^d = \prod_{k=1}^n \frac{1}{1 - \lambda_k x} = \frac{1}{x^n} \prod_{k=1}^n \frac{1}{x^{-1} - \lambda_k} = \frac{1}{x^n p_A(1/x)};$$

the apparent singularity at the origin is removable. Now observe that

$$\prod_{k=1}^n \frac{1}{1 - \lambda_k x} = \frac{1}{\det \operatorname{diag}(1 - \lambda_1 x, 1 - \lambda_2 x, \dots, 1 - \lambda_n x)} = \frac{1}{\det(I - xA)}$$

by the spectral theorem.  $\square$

**Example 20.** Let  $A = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}$ . Then  $p_A(z) = z^2 - z - 1$  and

$$\frac{1}{x^2 p_A(1/x)} = \frac{1}{1 - x - x^2} = \sum_{n=0}^{\infty} f_{n+1} x^n,$$

in which  $f_n$  is the  $n$ th Fibonacci number; these are defined by  $f_{n+2} = f_{n+1} + f_n$  and  $f_0 = 0$  and  $f_1 = 1$ . Thus,  $\|A\|_d^d = f_d$  for even  $d \geq 2$ .

**Remark 21.** If  $A \in H_n(\mathbb{C})$  is fixed, the sequence  $h_d(\lambda_1, \lambda_2, \dots, \lambda_n)$  satisfies a constant-coefficient recurrence of order  $n$  since its generating function is a rational function whose denominator has degree  $n$ . Solving such a recurrence is elementary, so one can compute  $\|A\|_d$  for  $d = 2, 4, 6, \dots$  via this method.

**Remark 22.** For small  $d$ , there is a simpler method. Since  $p_A(x)$  is monic, it follows that  $\widetilde{p}_A(x) = x^n p_A(1/x)$  has constant term 1. For small  $x$ , we have

$$\sum_{d=0}^{\infty} h_d(\lambda_1, \lambda_2, \dots, \lambda_n) x^d = \frac{1}{\widetilde{p}_A(x)} = \frac{1}{1 - (1 - \widetilde{p}_A(x))} = \sum_{d=0}^{\infty} (1 - \widetilde{p}_A(x))^d$$

so the desired  $h_d(\lambda_1, \lambda_2, \dots, \lambda_n)$  can be computed by the expanding the geometric series to the appropriate degree.

**Remark 23.** For  $d \geq 1$ , the Newton–Gerard identities imply

$$h_d(x_1, x_2, \dots, x_n) = \frac{1}{d} \sum_{i=1}^d h_{d-i}(x_1, x_2, \dots, x_n) p_i(x_1, x_2, \dots, x_n);$$

see [26, §10.12]. For  $A \in H_n(\mathbb{C})$  and  $d \geq 2$  even, it follows that

$$h_d(\lambda(A)) = \frac{1}{d} \sum_{i=1}^d h_{d-i}(\lambda(A)) \operatorname{tr}(A^i),$$

which can be used to compute  $\|A\|_d^d = h_d(\lambda(A))$  recursively.

**Remark 24.** If  $H, K \in H_n(\mathbb{C})$ , then  $\det(I - xH) = \det(I - xK)$  if and only if they are unitarily similar. However,  $H = \operatorname{diag}(1, 0)$  and  $K = \operatorname{diag}(1, -1)$  give

$$\frac{1}{\det(I - xH)} = \frac{1}{1 - x} = \sum_{j=0}^{\infty} x^j \quad \text{and} \quad \frac{1}{\det(I - xK)} = \frac{1}{1 - x^2} = \sum_{k=0}^{\infty} x^{2k},$$

so  $\|H\|_d = \|K\|_d$  for even  $d \geq 2$ . Of course, the odd-indexed coefficients (the complete homogeneous symmetric polynomials of odd degree) do not agree.

**4.2. Determinantal interpretation.** The material of the previous subsection leads to the determinantal interpretation (Theorem 6) stated in the introduction. We restate (and prove) the result here for convenience:

**Theorem 25.** *Let  $A \in M_n(\mathbb{C})$ . For  $d$  even,  $\binom{d}{d/2} \|A\|_d^d$  is the coefficient of  $z^{d/2} \bar{z}^{d/2}$  in the Taylor expansion of  $\det(I - zA - \bar{z}A^*)^{-1}$  about the origin.*

*Proof.* If  $H \in H_n(\mathbb{C})$ , the coefficient of  $x^d$  in  $\det(I - xH)^{-1}$  is  $\|H\|_d^d$  by Theorem 19. By plugging in  $H = zA + \bar{z}A^*$  and treating the resulting expression as a series in  $z$  and  $\bar{z}$ , Remark 18 implies that the coefficient of  $z^{d/2} \bar{z}^{d/2}$  equals  $\binom{d}{d/2} \|A\|_d^d$ .  $\square$

**Example 26.** Let  $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ . Then

$$\det(I - zA - \bar{z}A^*)^{-1} = \frac{1}{1 - \bar{z}z} = \sum_{n=0}^{\infty} \bar{z}^n z^n,$$

and hence  $\|A\|_d^d = \binom{d}{d/2}^{-1}$  for even  $d \geq 2$ .

**Example 27.** For

$$A = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \quad \text{we have} \quad \det(I - zA - \bar{z}A^*)^{-1} = \frac{1}{1 - z^3 - 3z\bar{z} - \bar{z}^3}.$$

Computer algebra reveals that  $\|A\|_2^2 = \|A\|_4^4 = \frac{3}{2}$ ,  $\|A\|_6^6 = \frac{29}{20}$ , and  $\|A\|_8^8 = \frac{99}{70}$ .

**Example 28.** The matrices

$$A = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & i \\ 0 & i & -1 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 0 & i \\ 1 & i & 0 \end{bmatrix}$$

satisfy

$$\det(I - zA - \bar{z}A^*)^{-1} = \frac{1}{1 - 4z\bar{z}} = \det(I - zB - \bar{z}B^*)^{-1},$$

so  $\|A\|_d = \|B\|_d$  for even  $d \geq 2$ . These matrices are not similar (let alone unitarily similar) since  $A$  is nilpotent of order two and  $B$  is nilpotent of order three.

**Remark 29.** In terms of the Laplace operator  $\Delta = \frac{\partial^2}{\partial z \partial \bar{z}}$ , Theorem 6 states that for even  $d$ ,

$$d! \|A\|_d^d = \Delta^{d/2} \frac{1}{\det(I - zA - \bar{z}A^*)} (0).$$

**4.3. Inner products.** Theorem 3 says that  $\|\cdot\|_d$  is a norm on  $M_n(\mathbb{C})$  for even  $d \geq 2$ . It is natural to ask when these norms are induced by an inner product.

**Theorem 30.** *The norm  $\|\cdot\|_d$  on  $M_n(\mathbb{C})$  (and its restriction to  $H_n(\mathbb{C})$ ) is induced by an inner product if and only if  $d = 2$  or  $n = 1$*

*Proof.* If  $n = 1$  and  $d \geq 2$  is even, then  $\|A\|_d$  is a fixed positive multiple of  $|a|$  for each  $A = [a] \in M_1(\mathbb{C})$ . Thus,  $\|\cdot\|_d$  on  $M_1(\mathbb{C})$  is induced by a positive multiple of the inner product  $\langle A, B \rangle = \bar{b}a$ , in which  $A = [a]$  and  $B = [b]$ .

If  $d = 2$  and  $n \geq 1$ , then  $\|A\|_2^2 = \frac{1}{2} \operatorname{tr}(A^*A) + \frac{1}{2} (\operatorname{tr} A) \operatorname{tr}(A^*)$ , which is induced by the inner product  $\langle A, B \rangle = \frac{1}{2} \operatorname{tr}(B^*A) + \frac{1}{2} (\operatorname{tr} B^*) (\operatorname{tr} A)$  on  $M_n(\mathbb{C})$ .

It suffices to show that in all other cases the norm  $\|A\|_d = (h_d(\lambda(A)))^{1/d}$  on  $H_n(\mathbb{R})$  does not arise from an inner product. For  $n \geq 2$ , let  $A = \text{diag}(1, 0, 0, \dots)$  and  $B = \text{diag}(0, 1, 0, \dots, 0) \in H_n(\mathbb{R})$ . Then  $\|A\|_d^2 = \|B\|_d^2 = 1$ . Next observe that  $\|A + B\|_d^2 = (d + 1)^{2/d}$  since there are exactly  $d + 1$  nonzero summands, each equal to 1, in the evaluation of  $h_d(\lambda(A + B))$ . Because of cancellation, a similar argument shows that  $\|A - B\|_d^2 = 1$ . A result of Jordan and von Neumann says that a vector space norm  $\|\cdot\|$  arises from an inner product if and only if it satisfies the parallelogram identity  $\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2)$  for all  $x, y$  [11]. If  $\|\cdot\|_d$  satisfies the parallelogram identity, then  $(d + 1)^{2/d} + 1 = 2(1 + 1)$ ; that is,  $(d + 1)^2 = 3^d$ . The solutions are  $d = 0$  (which does not yield an inner product) and  $d = 2$  (which, as we showed above, does). Thus, for  $n \geq 2$  and  $d \geq 2$ , the norm  $\|\cdot\|_d$  on  $H_n(\mathbb{C})$  does not arise from an inner product.  $\square$

**4.4. A tracial lower bound.** Each CHS norm on  $M_n(\mathbb{C})$  is bounded below by an explicit positive multiple of the trace seminorm. That is, the CHS norms of a matrix can be related to its mean eigenvalue.

**Theorem 31.** *For  $A \in M_n(\mathbb{C})$  and  $d \geq 2$  even,*

$$\|A\|_d \geq \binom{n+d-1}{d}^{1/d} \frac{\text{tr } A}{n}$$

*with equality if and only if  $A$  is a multiple of the identity.*

*Proof.* Let  $d \geq 2$  be even. For  $x = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ , let  $\tilde{x} = (\tilde{x}_1, \tilde{x}_2, \dots, \tilde{x}_n)$  denote its decreasing rearrangement (the notation  $x^\downarrow$  is frequently used in the literature). Then  $x$  majorizes  $y$ , denoted  $x \succeq y$ , if

$$\sum_{i=1}^k \tilde{x}_i \geq \sum_{i=1}^k \tilde{y}_i \quad \text{for } k = 1, 2, \dots, n, \quad \text{and} \quad \sum_{i=1}^n x_i = \sum_{i=1}^n y_i.$$

The even-degree complete homogeneous symmetric polynomials are Schur convex [29, Thm. 1]. That is,  $h_d(x) \geq h_d(y)$  whenever  $x \succeq y$ , with equality if and only if  $x$  is a permutation of  $y$ .

Let  $A \in M_n(\mathbb{C})$  and define  $B(t) = e^{it}A + e^{-it}A^*$  for  $t \in \mathbb{R}$ . Then  $\lambda(B(t))$  majorizes  $\mu(t) = (\mu(t), \mu(t), \dots, \mu(t)) \in \mathbb{R}^n$ , in which  $\mu(t) = \text{tr } B(t)/n$ . Thus,

$$\|B(t)\|_d^d = h_d(\lambda(B(t))) \geq h_d(\mu(t)) = \mu(t)^d \binom{n+d-1}{d}$$

with equality if and only if  $B(t) = \mu(t)I$ . It follows from (16) that

$$\|A\|_d \geq \left( \frac{\binom{n+d-1}{d}}{2\pi \binom{d}{d/2}} \int_0^{2\pi} \mu(t)^d dt \right)^{1/d}. \quad (17)$$

Combine this with

$$\begin{aligned} \int_0^{2\pi} \mu(t)^d dt &= \int_0^{2\pi} \left( \frac{\text{tr } B(t)}{n} \right)^d dt = \frac{1}{n^d} \int_0^{2\pi} (e^{it} \text{tr } A + e^{-it} \text{tr } A^*)^d dt \\ &= \frac{1}{n^d} \sum_{k=0}^d \binom{d}{k} (\text{tr } A^*)^{d-k} (\text{tr } A)^k \int_0^{2\pi} e^{i(2k-d)t} dt \end{aligned}$$

$$= \frac{2\pi}{n^d} \binom{d}{d/2} |\operatorname{tr} A|^d$$

and get the desired inequality. The continuity of the integrand ensures that equality occurs in (17) if and only if  $B(t) = \mu(t)I$  for all  $t \in \mathbb{R}$ . An operator-valued Fourier expansion reveals that  $e^{it}A + e^{-it}A^* = (\sum_{n \in \mathbb{Z}} \hat{\mu}(n)e^{int})I$ , so  $A = \hat{\mu}(1)I$ . Conversely, equality holds in (17) if  $A$  is a multiple of the identity.  $\square$

**Remark 32.** For each fixed  $n \geq 1$ , the constant  $\binom{n+d-1}{d}^{1/d}$  in Theorem 31 tends to 1 from above as  $d \rightarrow \infty$ . Therefore,  $\|A\|_d \geq \frac{1}{n} |\operatorname{tr} A|$  for all  $A \in M_n(\mathbb{C})$ .

**4.5. Monotonicity.** The next result shows how CHS norms relate to each other. For Hermitian matrices, the first inequality below is superior to the second.

**Theorem 33.** *Let  $2 \leq p < q$  be even.*

(a) *If  $A \in H_n(\mathbb{C})$ , then  $(p!)^{1/p} \|A\|_p \leq (q!)^{1/q} \|A\|_q$ .*

(b) *If  $A \in M_n(\mathbb{C})$ , then  $((\binom{p}{p/2})p!)^{1/p} \|A\|_p \leq ((\binom{q}{q/2})q!)^{1/q} \|A\|_q$ .*

*Proof.* (a) Let  $A \in H_n(\mathbb{C})$  have eigenvalues  $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n)$ , listed in decreasing order, and let  $\xi = (\xi_1, \xi_2, \dots, \xi_n)$  be a random vector, in which  $\xi_1, \xi_2, \dots, \xi_n$  are independent standard exponential random variables. Let  $d \geq 2$  be even and consider the random variable  $X = \langle \xi, \lambda \rangle$ . Then (10) ensures that

$$(d!)^{1/d} \|A\|_d = (d! h_d(\lambda))^{1/d} = \mathbb{E}[|\langle \xi, \lambda \rangle|^d]^{1/d} = \mathbb{E}[|X|^d]^{1/d} = \|X\|_{L^d}.$$

Since we are in a probability space (in particular, a finite measure space),  $\|X\|_{L^p} \leq \|X\|_{L^q}$  for  $0 < p < q < \infty$ . For  $2 \leq p < q$  even, this yields the desired inequality.

(b) Let  $A \in M_n(\mathbb{C})$  and let  $2 \leq p < q$  be even. For  $t \in [0, 2\pi]$ , (a) ensures that

$$\|e^{it}A + e^{-it}A^*\|_p^p \leq \frac{(q!)^{p/q}}{p!} \|e^{it}A + e^{-it}A^*\|_q^p.$$

Consider  $f(t) = \|e^{it}A + e^{-it}A^*\|_q$  as an element of  $L^p[0, 2\pi]$ . Hölder's inequality and (16) imply the desired inequality:

$$\begin{aligned} \|A\|_p &= \left( \frac{1}{2\pi \binom{p}{p/2}} \int_0^{2\pi} \|e^{it}A + e^{-it}A^*\|_p^p dt \right)^{1/p} \\ &\leq \left( \frac{1}{2\pi \binom{p}{p/2}} \right)^{1/p} \left( \frac{(q!)^{p/q}}{p!} \int_0^{2\pi} \|e^{it}A + e^{-it}A^*\|_q^p dt \right)^{1/p} \\ &\leq \frac{(q!)^{1/q}}{(p!)^{1/p}} \left( \frac{1}{2\pi \binom{p}{p/2}} \right)^{1/p} \|f\|_{L^p} \leq \frac{(q!)^{1/q}}{(p!)^{1/p}} \left( \frac{1}{2\pi \binom{p}{p/2}} \right)^{1/p} (2\pi)^{\frac{1}{p}-\frac{1}{q}} \|f\|_{L^q} \\ &\leq \frac{(q!)^{1/q}}{(p!)^{1/p}} \left( \frac{1}{2\pi \binom{p}{p/2}} \right)^{1/p} (2\pi)^{\frac{1}{p}-\frac{1}{q}} \left( \int_0^{2\pi} \|e^{it}A + e^{-it}A^*\|_q^q dt \right)^{1/q} \\ &\leq \frac{(q!)^{1/q}}{(p!)^{1/p}} \left( \frac{1}{\binom{p}{p/2}} \right)^{1/p} (2\pi)^{-\frac{1}{q}} \left( 2\pi \binom{q}{q/2} \right)^{1/q} \|A\|_q^q \end{aligned}$$

$$\leq \frac{((\binom{q}{q/2}q!)^{1/q}}{((\binom{p}{p/2}p!)^{1/p}} \|A\|_q^q. \quad \square$$

**Remark 34.** The previous result suggests that suitable constant multiples of the CHS norms may be preferable in some circumstances. However, the benefits appear to be outweighed by the cumbersome nature of these constants.

**Remark 35.** For  $A, B \in M_n(\mathbb{C})$ ,

$$\begin{aligned} 2\|AB\|_2^2 &= \operatorname{tr}(AB) \operatorname{tr}((AB)^*) + \operatorname{tr}((AB)^*AB) \\ &\leq 2 \operatorname{tr}(A^*A) \operatorname{tr}(B^*B) \\ &\leq 2(\operatorname{tr}(A) \operatorname{tr}(A^*) + \operatorname{tr}(A^*A))(\operatorname{tr}(B) \operatorname{tr}(B^*) + \operatorname{tr}(B^*B)) \\ &= 8\|A\|_2^2\|B\|_2^2, \end{aligned}$$

so  $2\|\cdot\|_2$  is submultiplicative. Actually, 2 is the smallest constant independent of  $n$  with this property, since

$$J = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$$

satisfies  $\|JJ^*\|_2 = 1 = 2\|J\|_2\|J^*\|_2$ .

**4.6. Symmetric Tensor Powers.** Let  $\mathcal{V}$  denote an  $n$ -dimensional  $\mathbb{R}$ -inner product space with orthonormal basis  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ . The  $k$ th tensor power of  $\mathcal{V}$  is the  $n^k$ -dimensional  $\mathbb{R}$ -inner product space  $\mathcal{V}^{\otimes k}$  spanned by the *simple tensors*

$$\mathbf{v}_{i_1} \otimes \mathbf{v}_{i_2} \otimes \cdots \otimes \mathbf{v}_{i_k}. \quad (18)$$

These simple tensors form an orthonormal basis of  $\mathcal{V}^{\otimes k}$ . An operator  $A : \mathcal{V} \rightarrow \mathcal{V}$  lifts to an operator on  $\mathcal{V}^{\otimes k}$  as follows. Define

$$A^{\otimes k}(\mathbf{v}_{i_1} \otimes \mathbf{v}_{i_2} \otimes \cdots \otimes \mathbf{v}_{i_k}) = A\mathbf{v}_{i_1} \otimes A\mathbf{v}_{i_2} \otimes \cdots \otimes A\mathbf{v}_{i_k}$$

and use the linearity of  $A$  and  $\otimes$  to write this in terms of the basis vectors (18). An important fact is that  $\mathcal{V}^{\otimes k}$  is basis-independent, in the sense of abstract nonsense.

The  $k$ th *symmetric tensor power* of  $\mathcal{V}$  is the  $\binom{n+k-1}{k}$ -dimensional vector space  $\operatorname{Sym}_k \mathcal{V} \subset \mathcal{V}^{\otimes k}$  spanned by the *symmetric tensors*:

$$\mathbf{v}_{i_1} \odot \mathbf{v}_{i_2} \odot \cdots \odot \mathbf{v}_{i_k} = \frac{1}{k!} \sum_{\sigma \in S_k} \mathbf{v}_{\sigma(i_1)} \otimes \mathbf{v}_{\sigma(i_2)} \otimes \cdots \otimes \mathbf{v}_{\sigma(i_k)},$$

where  $S_k$  denotes the symmetric group on  $k$  letters.

**Proposition 36.** *If  $d \geq 2$  is even and  $A \in H_n(\mathbb{C})$ , then*

$$\|A\|_d^d = \operatorname{tr}(A^{\operatorname{Sym}_d}).$$

*Proof.* Let  $A : \mathcal{V} \rightarrow \mathcal{V}$  be selfadjoint with eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$  and corresponding orthonormal eigenbasis  $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n$ . Define  $A^{\operatorname{Sym}_k} : \operatorname{Sym}_k \mathcal{V} \rightarrow \operatorname{Sym}_k \mathcal{V}$  by linear extension. Then  $\mathbf{v}_{i_1} \odot \mathbf{v}_{i_2} \odot \cdots \odot \mathbf{v}_{i_k}$  is an eigenvector of  $A^{\operatorname{Sym}_k}$  with eigenvalue  $\lambda_{i_1} \lambda_{i_2} \cdots \lambda_{i_k}$ . Sum over these  $\binom{n+k-1}{k}$  eigenvectors and conclude that  $\operatorname{tr}(A^{\operatorname{Sym}_k}) = h_k(\lambda_1, \lambda_2, \dots, \lambda_n)$ .  $\square$

If  $A$  is the adjacency matrix of a graph  $\Gamma$ , then  $\|A\|_d$  concerns the  $d$ th symmetric tensor power of  $\Gamma$ , a weighted graph obtained from  $\Gamma$  in a straightforward (but tedious) manner by computing the matrix representation of  $A^{\text{Sym}_d}$  with respect to the normalization of the orthogonal basis of symmetrized tensors.

**4.7. Equivalence constants.** Any two norms on a finite-dimensional vector space are equivalent. Thus, each norm  $\|\cdot\|_d$  on  $H_n(\mathbb{C})$  (with  $d \geq 2$  even) is equivalent to the operator norm  $\|\cdot\|_{\text{op}}$ . We compute admissible equivalence constants below.

**Theorem 37.** For  $A \in H_n(\mathbb{C})$  and even  $d \geq 2$ ,

$$\left(\frac{1}{2^{\frac{d}{2}}(\frac{d}{2})!}\right)^{1/d} \|A\|_{\text{op}} \leq \|A\|_d \leq \binom{n+d-1}{d}^{1/d} \|A\|_{\text{op}}$$

The upper inequality is sharp if and only if  $A$  is a multiple of the identity.

*Proof.* For  $A \in H_n(\mathbb{C})$  and even  $d \geq 2$ , the triangle inequality yields

$$\begin{aligned} \|A\|_d^d &= h_d(\lambda_1(A), \lambda_2(A), \dots, \lambda_n(A)) \\ &= |h_d(\lambda_1(A), \lambda_2(A), \dots, \lambda_n(A))| \\ &\leq h_d(|\lambda_1(A)|, |\lambda_2(A)|, \dots, |\lambda_n(A)|) \\ &\leq h_d(\|A\|_{\text{op}}, \|A\|_{\text{op}}, \dots, \|A\|_{\text{op}}) \\ &= \|A\|_{\text{op}}^d h_d(1, 1, \dots, 1) \\ &= \|A\|_{\text{op}}^d \binom{n+d-1}{d}. \end{aligned}$$

Equality occurs if and only if  $\lambda_i(A) = |\lambda_i(A)| = \|A\|_{\text{op}}$  for  $1 \leq i \leq n$ ; that is, if and only if  $A$  is a multiple of the identity.

Hunter [10] established that

$$h_{2p}(\mathbf{x}) \geq \frac{1}{2^p p!} \|\mathbf{x}\|^{2p},$$

in which  $\|\mathbf{x}\|$  denotes the Euclidean norm of  $\mathbf{x} \in \mathbb{R}^n$ . Let  $d = 2p$  and conclude

$$\|A\|_d \geq \left(\frac{1}{2^{\frac{d}{2}}(\frac{d}{2})!}\right)^{1/d} \|A\|_F \geq \left(\frac{1}{2^{\frac{d}{2}}(\frac{d}{2})!}\right)^{1/d} \|A\|_{\text{op}},$$

in which  $\|A\|_F$  denotes the Frobenius norm of  $A \in H_n(\mathbb{C})$ .  $\square$

**Remark 38.** For  $A \in M_n(\mathbb{C})$ , we may apply the upper bound in Theorem 37 to  $e^{it}A + e^{-it}A^*$  and use (14) to deduce that

$$\|A\|_d \leq \left(\frac{\binom{n+d-1}{d}}{2\pi(\frac{d}{2})} \int_0^{2\pi} \|e^{it}A + e^{-it}A^*\|_{\text{op}}^d dt\right)^{1/d} \leq 2 \left(\frac{\binom{n+d-1}{d}}{(\frac{d}{2})}\right)^{1/d} \|A\|_{\text{op}}.$$

**Remark 39.** Hunter's lower bound was improved by Baston [2], who proved that

$$h_{2p}(\mathbf{x}) \geq \frac{1}{2^p p!} \left(\sum_{i=1}^n x_i^2\right)^p + \lambda_p \left(\sum_{i=1}^n x_i\right)^{2p}$$

for  $\mathbf{x} = (x_1, x_2, \dots, x_n) \in \mathbb{R}^n$ , where

$$\lambda_p = \frac{1}{n^p} \left( \binom{n+2p-1}{2p} \frac{1}{n^p} - \frac{1}{2^p p!} \right) > 0.$$

Equality holds if and only if  $p = 1$  or  $p \geq 2$  and all the  $x_i$  are equal. However, Baston's result does not appear to yield a significant improvement in the lower bound of Theorem 37.

## 5. OPEN QUESTIONS

The answers to the following questions have eluded us.

**Problem 1.** *What are the best constants  $c_d$ , independent of  $n$ , such that  $c_d \|\cdot\|_d$  is submultiplicative? Do such constants exist? See Remark 35.*

**Problem 2.** *What is the best complexified version of Theorem 37? Can the upper bound be improved (the estimate  $\|e^{it}A + e^{-it}A^*\|_{\text{op}} \leq 2\|A\|_{\text{op}}$  seems wasteful on average)? Can we get a sharp lower bound?*

**Problem 3.** *If one uses (6) to evaluate  $\|A\|_d^d$ , there are many repeated terms. For example,  $(\text{tr } A^*A)(\text{tr } A)(\text{tr } A^*) = (\text{tr } AA^*)(\text{tr } A^*)(\text{tr } A)$  because of the cyclic invariance of the trace and the commutativity of multiplication. If one chooses a single representative for each such class of expressions and simplifies, one gets expressions such as (7) and (8). Is there a combinatorial interpretation of the resulting coefficients?*

For motivation, the reader is invited to consider

$$\begin{aligned} \|A\|_6^6 = & \frac{1}{720} \left( (\text{tr } A)^3 \text{tr}(A^*)^3 + 3 \text{tr}(A) \text{tr}(A^*)^3 \text{tr}(A^2) \right. \\ & + 9(\text{tr } A)^2 \text{tr}(A^*)^2 \text{tr}(A^*A) + 9 \text{tr}(A^*)^2 \text{tr}(A^2) \text{tr}(A^*A) \\ & + 18 \text{tr}(A) \text{tr}(A^*) \text{tr}(A^*A)^2 + 6 \text{tr}(A^*A)^3 + 3(\text{tr } A)^3 \text{tr}(A^*) \text{tr}(A^{*2}) \\ & + 9 \text{tr}(A) \text{tr}(A^*) \text{tr}(A^2) \text{tr}(A^{*2}) + 9(\text{tr } A)^2 \text{tr}(A^*A) \text{tr}(A^{*2}) \\ & + 9 \text{tr}(A^2) \text{tr}(A^*A) \text{tr}(A^{*2}) + 2 \text{tr}(A^*)^3 \text{tr}(A^3) \\ & + 6 \text{tr}(A^*) \text{tr}(A^{*2}) \text{tr}(A^3) + 18 \text{tr}(A) \text{tr}(A^*)^2 \text{tr}(A^*A^2) \\ & + 36 \text{tr}(A^*) \text{tr}(A^*A) \text{tr}(A^*A^2) + 18 \text{tr}(A) \text{tr}(A^{*2}) \text{tr}(A^*A^2) \\ & + 18(\text{tr } A)^2 \text{tr}(A^*) \text{tr}(A^{*2}A) + 18 \text{tr}(A^*) \text{tr}(A^2) \text{tr}(A^{*2}A) \\ & + 36 \text{tr}(A) \text{tr}(A^*A) \text{tr}(A^{*2}A) + 36 \text{tr}(A^*A^2) \text{tr}(A^{*2}A) \\ & + 2(\text{tr } A)^3 \text{tr}(A^{*3}) + 6 \text{tr}(A) \text{tr}(A^2) \text{tr}(A^{*3}) \\ & + 4 \text{tr}(A^3) \text{tr}(A^{*3}) + 18 \text{tr}(A^*)^2 \text{tr}(A^*A^3) \\ & + 18 \text{tr}(A^{*2}) \text{tr}(A^*A^3) + 18 \text{tr}(A) \text{tr}(A^*) \text{tr}(A^*AA^*A) \\ & + 18 \text{tr}(A^*A) \text{tr}(A^*AA^*A) + 36 \text{tr}(A) \text{tr}(A^*) \text{tr}(A^{*2}A^2) \\ & + 36 \text{tr}(A^*A) \text{tr}(A^{*2}A^2) + 18(\text{tr } A)^2 \text{tr}(A^{*3}A) \\ & + 18 \text{tr}(A^2) \text{tr}(A^{*3}A) + 36 \text{tr}(A^*) \text{tr}(A^*AA^*A^2) \\ & + 36 \text{tr}(A^*) \text{tr}(A^{*2}A^3) + 36 \text{tr}(A) \text{tr}(A^{*2}AA^*A) \\ & + 36 \text{tr}(A) \text{tr}(A^{*3}A^2) + 12 \text{tr}(A^*AA^*AA^*A) + 36 \text{tr}(A^{*2}A^2A^*A) \\ & \left. + 36 \text{tr}(A^{*2}AA^*A^2) + 36 \text{tr}(A^{*3}A^3) \right). \end{aligned}$$



## REFERENCES

- [1] A. I. Barvinok, *Low rank approximations of symmetric polynomials and asymptotic counting of contingency tables*, <https://arxiv.org/abs/math/0503170>.
- [2] V. J. Baston, *Two inequalities for the complete symmetric functions*, Math. Proc. Cambridge Philos. Soc. **84** (1978), no. 1, 1–3. MR 485422
- [3] Patrick Billingsley, *Probability and measure*, Wiley Series in Probability and Statistics, John Wiley & Sons, Inc., Hoboken, NJ, 2012, Anniversary edition [of MR1324786], With a foreword by Steve Lalley and a brief biography of Billingsley by Steve Koppes. MR 2893652
- [4] Albrecht Böttcher, Stephan Ramon Garcia, Mohamed Omar, and Christopher O’Neill, *Weighted means of B-splines, positivity of divided differences, and complete homogeneous symmetric polynomials*, Linear Algebra Appl. **608** (2021), 68–83. MR 4140644
- [5] Eduardo Marques de Sá, *Exposed faces and duality for symmetric and unitarily invariant norms*, vol. 197/198, 1994, Second Conference of the International Linear Algebra Society (ILAS) (Lisbon, 1992), pp. 429–450. MR 1275626
- [6] Stephan Ramon Garcia, Mohamed Omar, Christopher O’Neill, and Samuel Yih, *Factorization length distribution for affine semigroups II: asymptotic behavior for numerical semigroups with arbitrarily many generators*, J. Combin. Theory Ser. A **178** (2021), 105358, 34. MR 4175889
- [7] G. H. Hardy, J. E. Littlewood, and G. Pólya, *Inequalities*, Cambridge Mathematical Library, Cambridge University Press, Cambridge, 1988, Reprint of the 1952 edition. MR 944909
- [8] J. William Helton and Victor Vinnikov, *Linear matrix inequality representation of sets*, Comm. Pure Appl. Math. **60** (2007), no. 5, 654–674. MR 2292953
- [9] Alfred Horn, *Eigenvalues of sums of Hermitian matrices*, Pacific J. Math. **12** (1962), 225–241. MR 140521
- [10] D. B. Hunter, *The positive-definiteness of the complete symmetric functions of even order*, Math. Proc. Cambridge Philos. Soc. **82** (1977), no. 2, 255–258. MR 450079
- [11] P. Jordan and J. Von Neumann, *On inner products in linear, metric spaces*, Ann. of Math. (2) **36** (1935), no. 3, 719–723. MR 1503247
- [12] Igor Klep, Špela Špenko, and Jurij Volčič, *Positive trace polynomials and the universal Procesi-Schacher conjecture*, Proc. Lond. Math. Soc. (3) **117** (2018), no. 6, 1101–1134. MR 3893175
- [13] Alexander A. Klyachko, *Stable bundles, representation theory and Hermitian operators*, Selecta Math. (N.S.) **4** (1998), no. 3, 419–445. MR 1654578
- [14] Allen Knutson and Terence Tao, *The honeycomb model of  $GL_n(\mathbb{C})$  tensor products. I. Proof of the saturation conjecture*, J. Amer. Math. Soc. **12** (1999), no. 4, 1055–1090. MR 1671451
- [15] A. S. Lewis, *Convex analysis on the Hermitian matrices*, SIAM J. Optim. **6** (1996), no. 1, 164–177. MR 1377729
- [16] ———, *Group invariance and convex matrix analysis*, SIAM J. Matrix Anal. Appl. **17** (1996), no. 4, 927–949. MR 1410709
- [17] M. Marcus, *An eigenvalue inequality for the product of normal matrices*, Amer. Math. Monthly **63** (1956), 173–174. MR 75920
- [18] Albert W. Marshall, Ingram Olkin, and Barry C. Arnold, *Inequalities: theory of majorization and its applications*, second ed., Springer Series in Statistics, Springer, New York, 2011. MR 2759813
- [19] J. B. McLeod, *On four inequalities in symmetric functions*, Proc. Edinburgh Math. Soc. **11** (1958/1959), 211–219. MR 0112935
- [20] Leon Mirsky, *On the trace of matrix products*, Math. Nachr. **20** (1959), 171–174. MR 125851
- [21] Gustavo A. Muñoz, Yannis Sarantopoulos, and Andrew Tonge, *Complexifications of real Banach spaces, polynomials and multilinear maps*, Studia Math. **134** (1999), no. 1, 1–33. MR 1688213
- [22] Franz Rendl and Henry Wolkowicz, *Applications of parametric programming and eigenvalue maximization to the quadratic assignment problem*, Math. Programming **53** (1992), no. 1, Ser. A, 63–78. MR 1151765
- [23] James Renegar, *Hyperbolic programs, and their derivative relaxations*, Found. Comput. Math. **6** (2006), no. 1, 59–79. MR 2198215
- [24] Hans Richter, *Zur Abschätzung von Matrizennormen*, Math. Nachr. **18** (1958), 178–187. MR 111758
- [25] Ionel Roventă and Laurențiu Emanuel Temereancă, *A note on the positivity of the even degree complete homogeneous symmetric polynomials*, Mediterr. J. Math. **16** (2019), no. 1, Paper No. 1, 16. MR 3887204
- [26] Raymond Séroul, *Programming for mathematicians*, Universitext, Springer-Verlag, Berlin, 2000, Translated from the 1995 French original by Donal O’Shea. MR 1740388

- [27] Suvrit Sra, *New concavity and convexity results for symmetric polynomials and their ratios*, *Linear Multilinear Algebra* **68** (2020), no. 5, 1031–1038. MR 4121426
- [28] Richard P. Stanley, *Enumerative combinatorics. Vol. 2*, *Cambridge Studies in Advanced Mathematics*, vol. 62, Cambridge University Press, Cambridge, 1999, With a foreword by Gian-Carlo Rota and appendix 1 by Sergey Fomin. MR 1676282
- [29] Terence Tao, *Schur convexity and positive definiteness of the even degree complete homogeneous symmetric polynomials*, <https://terrytao.wordpress.com/2017/08/06/schur-convexity-and-positive-definiteness-of-the-even-degree-complete-homogeneous-symmetric-polynomials/>.
- [30] C. M. Theobald, *An inequality for the trace of the product of two symmetric matrices*, *Math. Proc. Cambridge Philos. Soc.* **77** (1975), 265–267. MR 414593
- [31] John von Neumann, *“Some matrix inequalities” in Collected Works. Vol. IV: Continuous geometry and other topics*, Pergamon Press, Oxford-London-New York-Paris, 1962, General editor: A. H. Taub. MR 0157874
- [32] David G. Wagner, *Multivariate stable polynomials: theory and applications*, *Bull. Amer. Math. Soc. (N.S.)* **48** (2011), no. 1, 53–84. MR 2738906
- [33] J. N. Whiteley, *Some inequalities concerning symmetric forms*, *Mathematika* **5** (1958), 49–57. MR 95233

DEPARTMENT OF MATHEMATICS, POMONA COLLEGE, 610 N. COLLEGE AVE., CLAREMONT, CA 91711

Email address: [konrad.aguilar@pomona.edu](mailto:konrad.aguilar@pomona.edu)

URL: <https://aguilar.sites.pomona.edu/>

Email address: [Angel.Chavez@pomona.edu](mailto:Angel.Chavez@pomona.edu)

Email address: [stephan.garcia@pomona.edu](mailto:stephan.garcia@pomona.edu)

URL: <http://pages.pomona.edu/~sg064747>

DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, TAMU 3368, COLLEGE STATION, TX 77843-3368

Email address: [volcic@tamu.edu](mailto:volcic@tamu.edu)